

DEFINITION

- Complains of upper, mid, or lower back pain that occurs mainly in the midline.
- Not due to a traumatic injury
- Minor muscle strain and overuse are covered in this protocol. Sciatic pain is also covered.

PAIN SEVERITY is defined as:

- MILD (1-3): doesn't interfere with normal activities
- MODERATE (4-7): interferes with normal activities or awakens from sleep
- SEVERE (8-10): excruciating pain, unable to do any normal activities

Notes:

- Pain in the back from significant blunt or penetrating trauma should be triaged using the **Back Injury** protocol.
- Pain in the lower back in pregnant women, consider labor. See **Pregnancy - Back Pain**, **Pregnancy - Labor**, and **Pregnancy - Labor, Preterm** protocol.

TRIAGE ASSESSMENT QUESTIONS

Call EMS 911 Now

Passed out (i.e., lost consciousness, collapsed and was not responding)

R/O: ruptured abdominal aortic aneurysm. FIRST AID: Lie down with the feet elevated.

Shock suspected (e.g., cold/pale/clammy skin, too weak to stand, low BP, rapid pulse)

R/O: ruptured abdominal aortic aneurysm. FIRST AID: Lie down with the feet elevated.

Sounds like a life-threatening emergency to the triager

See More Appropriate Protocol

Major injury to the back (e.g., MVA, fall > 10 feet or 3 meters, penetrating injury, etc.)

Go to Protocol: Back Injury (Adult)

Pain in the upper back over the ribs (rib cage) that radiates (travels) into the chest

Go to Protocol: Chest Pain (Adult)

Pain in the upper back over the ribs (rib cage) and worsened by coughing (or clearly increases with breathing)

Go to Protocol: Chest Pain (Adult)

Back pain during pregnancy

Go to Protocol: Pregnancy - Back Pain (Adult)

Go to ED Now

Severe back pain of sudden onset and age > 60 years

R/O: compression fracture, abdominal aortic aneurysm

Severe abdominal pain (e.g., excruciating)

Abdominal pain and age > 60 years

R/O: compression fracture, aortic aneurysm

Unable to urinate (or only a few drops) and bladder feels very full

R/O: urinary retention, cauda equina syndrome, conus medullaris syndrome

Loss of bladder or bowel control (urine or bowel incontinence; wetting self, leaking stool) of new-onset

R/O: urinary retention with overflow incontinence, cauda equina syndrome, conus medullaris syndrome

Numbness (loss of sensation) in groin or rectal area

R/O: cauda equina syndrome, conus medullaris syndrome

Go to ED/UCC Now (or to Office with PCP Approval)

Pain radiates into groin, scrotum

R/O: kidney stones

Blood in urine (red, pink, or tea-colored)

Vomiting and pain over lower ribs of back (i.e., flank - kidney area)

Weakness of a leg or foot (e.g., unable to bear weight, dragging foot)

R/O: nerve root impingement or cord compression

Patient sounds very sick or weak to the triager

Reason: Severe acute illness or serious complication suspected.

Go to Office Now

Fever > 100.4° F (38.0° C) and flank pain

R/O: pyelonephritis

Pain or burning with passing urine (urination)

R/O: pyelonephritis

See in Office Today

Severe back pain (e.g., excruciating, unable to do any normal activities) and not improved after pain medicine and **Care Advice**

Numbness in an arm or hand (i.e., loss of sensation) and upper back pain

R/O: cervical nerve root compression, herniated disk; with pain radiating into upper back

Numbness in a leg or foot (i.e., loss of sensation)

R/O: severe back strain, cord compression

High-risk adult (e.g., history of cancer, history of HIV, or history of IV Drug Use)

R/O: metastasis, epidural abscess

Soft tissue infection (e.g., abscess, cellulitis) or other serious infection (e.g., bacteremia) in last 2 weeks

Reason: Recent soft tissue or blood infection has higher risk of a spinal infection.

Painful rash with multiple small blisters grouped together (i.e., dermatomal distribution or "band" or "stripe")

R/O: herpes zoster (shingles)

Pain radiates into the thigh or further down the leg, and in both legs

Reason: Bilateral sciatica carries higher risk.

See in Office Today or Tomorrow

Age > 50 and no history of prior similar back pain

Reason: Higher risk of serious medical cause.

See in Office Within 3 Days

Moderate back pain (e.g., interferes with normal activities) and present > 3 days

R/O: sciatica

Pain radiates into the thigh or further down the leg

R/O: sciatica

Patient wants to be seen

See in Office Within 2 Weeks

Back pain lasts > 2 weeks

Back pain is a chronic symptom (recurrent or ongoing AND lasting > 4 weeks)

Home Care

Caused by a twisting, bending, or lifting injury

R/O: muscle strain, overuse

Caused by overuse from recent vigorous activity (e.g., exercise, gardening, lifting and carrying, sports)

R/O: muscle strain, overuse

Back pain

R/O: muscle strain, overuse

Preventing back strain, questions about

HOME CARE ADVICE

Back Pain

1. Reassurance and Education - Back Pain:

- Back pain can result from excessive twisting, heavy lifting, or from un-noticed minor back injuries.
- Usually the pain gets better in 1 to 2 weeks.

- You can treat most back pain at home.
 - *Here is some care advice that should help.*
2. **Cold or Heat:**
 - **Cold Pack:** For pain or swelling, use a cold pack or ice wrapped in a wet cloth. Put it on the sore area for 20 minutes. Repeat 4 times on the first day, then as needed.
 - **Heat Pack:** If pain lasts over 2 days, apply heat to the sore area. Use a heat pack, heating pad, or warm wet washcloth. Do this for 10 minutes, then as needed. For widespread stiffness, take a hot bath or hot shower instead. Move the sore area under the warm water.
 3. **Sleep:**
 - Sleep on your side with a pillow between your knees. If you sleep on your back, put a pillow under your knees.
 - Avoid sleeping on your stomach.
 - Your mattress should be firm. Avoid waterbeds.
 4. **Activity:**
 - Keep doing your day-to-day activities if it is not too painful. Staying active is better than resting.
 - Avoid anything that makes your pain worse. Avoid heavy lifting, twisting, and too much exercise until your back heals.
 - You do not need to stay in bed.
 5. **Pain Medicines:**
 - For pain relief, you can take either acetaminophen, ibuprofen, or naproxen.
 - They are over-the-counter (OTC) pain drugs. You can buy them at the drugstore.
 - **Acetaminophen - Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4 to 6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
 - **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
 - **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.
 - **Naproxen (e.g., Aleve):** Take 220 mg (one 220 mg pill) by mouth every 8 to 12 hours as needed. You may take 440 mg (two 220 mg pills) for your first dose. The most you should take each day is 660 mg (three 220 mg pills a day), unless your doctor has told you to take more.
 6. **Pain Medicines - Extra Notes and Warnings:**
 - Use the lowest amount of medicine that makes your pain better.
 - Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
 - McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, the maximum dose per day is ten Regular-Strength (325 mg) pills.
 - Bayer, the company that makes Aleve, has different dosage instructions for Aleve in Canada and the United States. In Canada, the maximum recommended dose per day is 440 mg (2 pills or caplets). In the United States, the maximum dose per day is 660 mg (3 pills or caplets).
 - **Caution:** Do not take acetaminophen if you have liver disease.
 - **Caution:** Do not take ibuprofen or naproxen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do

not take ibuprofen or naproxen for more than 7 days without consulting your doctor.

- *Before taking any medicine, read all the instructions on the package.*

7. **Call Back If:**

- Fever occurs
- Numbness or weakness occurs, or bowel/bladder problems
- Pain begins to shoot into the leg
- Pain persists over 2 weeks
- Pain becomes worse
- You become worse

Back Pain From Lifting or Twisting

1. **Reassurance and Education - Back Pain from Lifting or Twisting:**

- Twisting or heavy lifting can cause back pain. It can also occur after un-noticed minor back injuries
- With treatment, the pain most often goes away in 1 to 2 weeks.
- You can treat most back pain at home.
- *Here is some care advice that should help.*

2. **Use a Cold Pack for Pain:**

- Put a cold pack or an ice bag (wrapped in a moist towel) on the area for 20 minutes.
- Repeat in 1 hour, then every 4 hours while awake.
- Continue this for the first 48 hours (2 days).
- This will help decrease pain.
- *Caution:* Avoid frostbite.

3. **Use Heat After 48 Hours for Pain:**

- If pain lasts over 48 hours (2 days), put heat on the sore area.
- Use a heat pack, heating pad, or warm wet washcloth.
- Do this for 10 minutes three times a day.
- This will help increase blood flow and improve healing.
- *Caution:* Burn. Do not sleep on a heating pad.

4. **Sleep:**

- Sleep on your side with a pillow between your knees.
- If you sleep on your back, place a pillow under your knees.
- Avoid sleeping on the abdomen. The mattress should be firm or reinforced with a board.
- Avoid waterbeds.

5. **Activity:**

- Continue ordinary activities as much as your pain permits. Continued activity is more healing for the back than rest.
- Avoid any activities that cause severe pain.
- Use caution and commonsense when performing heavy lifting. Similarly, be careful during strenuous exercise.
- Note: Complete bed rest is unnecessary.

6. **Pain Medicines:**

- For pain relief, you can take either acetaminophen, ibuprofen, or naproxen.
- They are over-the-counter (OTC) pain drugs. You can buy them at the drugstore.
- **Acetaminophen - Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4 to 6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The

most you should take each day is 3,250 mg (10 pills a day).

- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).

- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

- **Naproxen (e.g., Aleve):** Take 220 mg (one 220 mg pill) by mouth every 8 to 12 hours as needed. You may take 440 mg (two 220 mg pills) for your first dose. The most you should take each day is 660 mg (three 220 mg pills a day), unless your doctor has told you to take more.

7. **Pain Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your pain better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, the maximum dose per day is ten Regular-Strength (325 mg) pills.
- Bayer, the company that makes Aleve, has different dosage instructions for Aleve in Canada and the United States. In Canada, the maximum recommended dose per day is 440 mg (2 pills or caplets). In the United States, the maximum dose per day is 660 mg (3 pills or caplets).
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen or naproxen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen or naproxen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

8. **Liniments:**

- If the patient presses for additional medication, a deep-heating cream has fewer side effects than muscle relaxants (e.g., drowsiness).

9. **Call Back If:**

- Fever occurs
- Numbness or weakness occurs, or bowel/bladder problems
- Pain begins to shoot into the leg
- Pain persists over 2 weeks
- Pain becomes worse
- You become worse

Preventing Back Strain

1. **Prevention:**

- The only way to prevent future backaches is to keep your back muscles in excellent physical condition.
- A sedentary lifestyle (lack of exercise) is a risk factor for developing back pain.
- Walking, stationary biking, and swimming provide good aerobic conditioning as well as exercise for your back.
- Being overweight puts more weight on the spine and thus increases the risk of back pain. If you are overweight, work with your doctor to develop a weight-loss program.

2. **Good Body Mechanics:**

- **Lifting:** Stand close to the object to be lifted. Keep your back straight and lift by bending your legs. Ask for lifting help if needed.
- **Sleeping:** Sleep on a firm mattress.
- **Sitting:** Avoid sitting for long periods of time without a break. Avoid slouching. Place a pillow or towel behind your lower back for support.
- **Posture:** Maintain good posture.

3. **Strengthening Exercises:**

- During the first couple days after an injury, strengthening exercises should be avoided. The following exercises can help strengthen the back. Perform the following exercises 3 to 10 times each day, for 5 to 10 seconds each time.
- **Bent knee sit-ups:** Lay on back, curl forward lifting shoulders about 6 inches (15 cm) off the floor.
- **Leg lifts:** Lay on back, lift foot 6 inches (15 cm) off floor (one leg at a time).
- **Pelvic tilt:** Lay on back with knees bent, push lower back against floor.
- **Chest lift:** Lie face down on ground, place arms by your sides, lift shoulders off the floor.

4. **Call Back If:**

- You have more questions
- You become worse

FIRST AID

FIRST AID Advice for Shock: Lie down with the feet elevated.

BACKGROUND INFORMATION

Key Points

- Lower back pain is a cause of countless visits to physicians' offices and emergency departments. It is the second most common cause of lost workdays, after cold and flu symptoms. Over 80% of people at some point in their lives have lower back pain.
- Nearly 1 in 3 adults have back pain during any three month period. Back pain occurs more often in older individuals.
- However, there is some good news. In most cases, the back pain is not serious and it has a self-limited course. Pain subsides within 4-6 weeks in 90% of individuals experiencing acute low back pain.

Four Categories of Back Pain

- **Potentially serious:** Examples include abdominal aortic aneurysm, neoplasm, osteomyelitis, epidural abscess, vertebral fracture, and neurologic emergencies (e.g., cauda equina syndrome).
- **Sciatica** (back pain with neurologic symptoms): There is radiation of the back pain (or buttock pain) into a lower extremity suggesting lumbosacral nerve root compression. There may be associated leg weakness, numbness, or paresthesias.
- **Non-specific back pain:** No neurologic symptoms. Examples include lumbar strain/sprain, degenerative osteoarthritis, lumbar disc disease, and fibromyalgia.
- **Referred back pain:** There are gastrointestinal causes like pancreatitis, biliary colic, and posterior gastric ulcer; genitourinary causes like renal colic, pyelonephritis, endometriosis, and ovarian cyst.

Lumbar Strain

- Acute lower back pain in the 18 to 50 year old age group is usually a symptom of strain of some of the 200 muscles in the back that allow us to stand upright.
- Often the triggering event is carrying something too heavy, lifting from an awkward position, bending too far backward or sideways, or overuse.
- Individuals with strained back muscles often note that the pain is increased by bending or twisting movements, relieved by assuming certain positions, and that the back muscles are tender.

Degenerative Osteoarthritis

- Degenerative osteoarthritis is a common cause of back pain in the older adult population.
- In uncomplicated osteoarthritis, individuals will complain of chronic midline back discomfort. Frequently, there is morning stiffness that improves as the day progresses.

Bed Rest and Overtreatment

- Complete bed rest is inconvenient and unnecessary in the majority of patients, including those who need to be examined by the physician. Complete bed rest should never be recommended over the telephone.
- Research has demonstrated that continuing ordinary activities within the limits permitted by pain results in a speedier recovery than rest [Malmivaara reference].

REFERENCES

1. Argoff CA. Spinal and radicular pain disorders. *Neurol Clin.* 1998;16(4):833-50.
2. Atlas SJ, Nardin RA. Evaluation and treatment of low back pain: an evidenced-based approach to clinical care. *Muscle & Nerve.* 2003;27(3):265-84.
3. Boonstra AM, Stewart RE, Köke AJ, Oosterwijk RF, Swaan JL, Schreurs KM, Schiphorst Preuper HR. Cut-Off Points for Mild, Moderate, and Severe Pain on the Numeric Rating Scale for Pain in Patients with Chronic Musculoskeletal Pain, etc. *Front Psychol.* 2016 Sep 30;7:1466.
4. Chou R, Huffman LH; American Pain Society; American College of Physicians. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians. *Clin. Practice Guideline. Ann Intern Med.* 2007 Oct 2;147(7):492-504.
5. Dahm KT, Brurberg KG, Jamtvedt G, Hagen KB. Advice to rest in bed versus advice to stay active for acute low-back pain and sciatica. *Cochrane Database Syst Rev.* 2010 Jun 16;(6):CD007612.
6. Dart RC, Kuffner EK, Rumack BH. Treatment of pain or fever with paracetamol (acetaminophen) in the alcoholic patient: a systematic review. *Am J Ther.* 2000 Mar;7(2):123-34.
7. Davis WT, April MD, Mehta S, Long B, Shroyer S. High risk clinical characteristics for pyogenic spinal infection in acute neck or back pain: Prospective cohort study. *Am J Emerg Med.* 2020;38(3):491-496.
8. Devereaux MW. Low back pain. *Prim Care.* 2004; 31(1): 33-51.

9. Deyo RA, Weinstein JN. Low back pain. *N Eng J Med*. 2001;344:363-9.
10. Divoll M, Abernethy DR, Ameer B, Greenblatt DJ. Acetaminophen kinetics in the elderly. *Clin Pharmacol Ther*. 1982 Feb;31(2):151-6.
11. Fontenelle LF, Sarti TD. Kidney Stones: Treatment and Prevention. *Am Fam Physician*. 2019 Apr 15;99(8):490-496.
12. French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ Superficial heat or cold for low back pain. *Cochrane Database Syst Rev*. 2006 Jan 25;(1):CD004750.
13. Friedman BW, Irizarry E, Solorzano C, Zias E, Pearlman S, Wollowitz A, Jones MP, Shah PD, Gallagher EJ. A Randomized, Placebo-Controlled Trial of Ibuprofen Plus Metaxalone, Tizanidine, or Baclofen for Acute Low Back Pain. *Ann Emerg Med*. 2019 Oct;74(4):512-520.
14. Griffin G, Tudiver F, Grant WD. Do NSAIDs help in acute or chronic low back pain? *Am Fam Physician*. 2002;65(7):1319-21.
15. Hilde G, Hagen KB, Jamtvedt G, Winnem M. Advice to stay active as a single treatment for low back pain and sciatica. *Cochrane Database Syst Rev*. 2002;(2):CD003632.
16. Hróbjartsson A, Gøtzsche PC. Placebo interventions for all clinical conditions. *Cochrane Database Syst Rev*. 2010 Jan 20;(1):CD003974.
17. Johnson JR, Russo TA. Acute Pyelonephritis in Adults. *N Engl J Med*. 2018 Jan 4;378(1):48-59.
18. Kovacs FM, Abaira V, Pena A: Effect of firmness of mattress on chronic non-specific low-back pain: randomised, double-blind, controlled, multicentre trial. *Lancet*. 2003 Nov 15;362(9396):1599-604.
19. Malmivaara A, Hakkinen U, Aro T, et al. The treatment of acute low back pain: bed rest, exercises, or ordinary activity? *N Engl J Med*. 1995;332:351-355.
20. Motov S, Masoudi A, Drapkin J, et al. Comparison of Oral Ibuprofen at Three Single-Dose Regimens for Treating Acute Pain in the Emergency Department: A Randomized Controlled Trial. *Ann Emerg Med*. 2019;74(4):530-537.
21. No authors listed. QuickStats: Percentage of Adults Aged ≥18 Years Who Had Lower Back Pain in the Past 3 Months, by Sex and Age Group - National Health Interview Survey, United States, 2018. *MMWR Morb Mortal Wkly Rep*. 2020;68(5152):1196.
22. No authors listed. QuickStats: Percentage of Emergency Department Visits for Pain at Which Opioids Were Given or Prescribed, by Geographic Region of the Hospital United States, 2005-2017 *MMWR Morb Mortal Wkly Rep*. 2018;67(50):1400.
23. Rainsford KD. Ibuprofen: pharmacology, efficacy and safety. *Inflammopharmacology*. 2009 Dec;17(6):275-342.
24. Roelofs PD, Deyo RA, Koes BW, Scholten RJ, van Tulder MW. Non-steroidal anti-inflammatory drugs for low back pain. *Cochrane Database Syst Rev*. 2008 Jan 23;(1):CD000396.
25. Sampson FC, Goodacre SW, O'Cathain A. The Reality of Pain Scoring in the Emergency Department: Findings From a Multiple Case Study Design. *Ann Emerg Med*. 2019 Oct;74(4):538-548. *Ann Emerg Med*. 2019 Oct;74(4):512-520.

26. Schieber LZ, Guy GP Jr, Seth P, Losby JL. Variation in Adult Outpatient Opioid Prescription Dispensing by Age and Sex - United States, 2008-2018. MMWR Morb Mortal Wkly Rep. 2020;69(11):298-302.
27. Swenson R Lower back pain; differential diagnosis: a reasonable clinical approach. Neurol Clin. 1999;17(1):43-63.
28. Takala E. Immediate referral from general practice for lumbar spine X-ray does not improve quality of life or pain for people with lower back pain. Evidence-based Healthcare. 2002; 6(2):91-92.

AUTHOR AND COPYRIGHT

Author:	David A. Thompson, MD, FACEP
Copyright:	2000-2022, LaGrange Medical Software, Inc. All rights reserved.
Company:	Schmitt-Thompson Clinical Content
Content Set:	Office Hours Telehealth Triage Protocols Adult
Version Year:	2022
Last Revised:	5/17/2022
Last Reviewed:	3/12/2022